



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2011


Effects of pasture type and animal breed on bone metabolism of lambs grazing high altitude grasslands

Willems, Helen ; Leiber, Florian ; Kreuzer, Michael ; Liesegang, Annette

Posted at the Zurich Open Repository and Archive, University of Zurich
ZORA URL: <https://doi.org/10.5167/uzh-73416>
Conference or Workshop Item


Originally published at:

Willems, Helen; Leiber, Florian; Kreuzer, Michael; Liesegang, Annette (2011). Effects of pasture type and animal breed on bone metabolism of lambs grazing high altitude grasslands. In: The 8th International Symposium on the Nutrition of Herbivores (ISNH8), Aberystwyth, Wales, September 2011.



Advances in **Animal Biosciences**

Proceedings of the 8th International
Symposium on the Nutrition of Herbivores
(ISNH8)



advances in
animal
biosciences

CAMBRIDGE
UNIVERSITY PRESS

Proceedings
of the 8th International Symposium on the
Nutrition of Herbivores
2011

Advances in Animal Biosciences

This book is part of a series which is a companion to the journal ANIMAL



The Stapledon Memorial Trust

Eighth International Symposium on the Nutrition of Herbivores

(ISNH8)



6th-9th September 2011, Aberystwyth, Wales, UK, www.isnh8.org

These Proceedings of the International Symposium on the Nutrition of Herbivores constitute summaries of papers presented at the 8th conference in Aberystwyth, Wales UK, 6th–9th September 2011.

The summaries have not been edited and the Organising Committee and publisher accept no responsibility for their accuracy. Views expressed in all contributions are those of the authors and not those of the symposium's committee members.

This publication contains all summaries that were available at the time of going to press.

ISNH8 COMMITTEE MEMBERS

Local Organising Committee

Nigel Scollan, Chair
Jamie Newbold
John Healey
Christina Marley
Michael Lee
Michael Rose
Neil McEwan
Jon Moorby
Siwan Griffiths
Roger Merry
Alison Kingston-Smith
Helen Lloyd
Jessica Longworth

International Advisory Committee

Hugh Dove, Chairperson, Australia
Andrew Sykes, New Zealand
René Baumont, France
Isabelle Cassar-Malek, France
Octavio Castelan-Ortega, Mexico
Jan Dijkstra, The Netherlands
Simone Hoskins, New Zealand
Denis Krause, Canada
Rod Mackie, USA
Ilias Kyriazakis, United Kingdom
Ruijun Long, China
Chris McSweeney, Australia
Qingxiang Meng, China
Bob Ørskov, United Kingdom
Dennis Poppi, Australia
Nigel Scollan, United Kingdom
Abeliza Kimambo, Tanzania
Peiqiang Yu, Canada

This publication has been produced with the kind support of the Stapledon Memorial Trust

© 2011 Animal Consortium
ISSN 2040-4700

PROGRAMME

THEATRE PRESENTATIONS

SESSION 1: Role of herbivores in sustainable land production systems

- 241 **The role of sheep in saltbush domestication - what can they tell us?**
Hayley Norman, Peter Jessop, Matt Wilmot
- 242 **Intensification alternative for pasture-based systems in Australia: a complementary forage system whole farm study**
Santiago Fariña, Sergio Garcia, William Fulkerson

SESSION 2: Nutrition and health of herbivores

- 243 **Effects of two protein sources on the periparturient relaxation of immunity to gastrointestinal nematode parasites**
Panagiotis Sakkas, Jos Houdijk, Spiridoula Ahanasiadou, Ilias Kyriazakis
- 244 **Crude glycerin as an alternative energy source for dairy cows in mid lactation**
Cassio Andre Wilbert, Enio Rosa Prates, Julio Otavio Barcellos, Jorge Schafhäuser
- 245 **Effects of irrigation and rates and timing of nitrogen fertiliser on proportions of maize plant fractions and nutritive value of whole crop maize silage for dairy cows**
M.R. Islam, S.C. Garcia
- 246 **Effects of dietary molybdenum, sulphur and zinc on the excretion and hepatic accumulation of copper in sheep fed palm kernel cake based diets**
Abdul Razak Alimon, Michael Ivan, Syed Jalaludin
- 247 **Mineral supplementation of lambing merino ewes grazing dual-purpose wheat under Australian conditions increases twin-lamb growth rate**
Shawn McGrath, Marie Bhanugopan, James Virgona, Hugh Dove, Michael Friend
- 248 **Effects of pasture type and animal breed on bone metabolism of lambs grazing high altitude grasslands**
Helen Willems, Florian Leiber, Michael Kreuzer, Annette Liesegang
- 249 **Evaluation of fresh lettuce wastes in comparison with fresh pawpaw leaves, whole lettuce and cabbage wastes as sole feed ingredient for growing *Archachatina marginata* snails**
Olubukola Babalola

SESSION 3: Novel developments in herbivore nutrition: molecular and cellular aspects

- 250 **Bacterial protein breakdown by different rumen protozoal groups**
Alejandro Belanche, Gabriel de la Fuente, Antonio I Martín-García, Jon M Moorby, Charles J Newbold

Effects of pasture type and animal breed on bone metabolism of lambs grazing high altitude grasslands

H Willems¹, F Leiber¹, M Kreuzer¹, A Liesegang²

¹ETH Zurich, Institute of Agricultural Sciences IAS, Zurich, Switzerland, ²University of Zurich, Institute of Animal Nutrition, Zurich, Switzerland

Email: hwillems@ethz.ch

Introduction High alpine grasslands serve as feed resource for ruminant livestock species during summer. Low feed density and the steep slopes in alpine regions require large movement activities of the grazing animal (Liesegang *et al.*, 2010). Due to their body size, sheep are well adapted to graze on those topographically and climatically harsh environments. Depending on magnitude and rate of strain, movement activities may influence bone metabolism in different ways (Bass *et al.*, 2005). The present study aimed at investigating the effects of grazing on different alpine pasture types, characterised by vegetation type, slope and Ca content, and animal breed on bone metabolism of lambs.

Material and methods Lambs of two Swiss mountain breeds, Engadin Sheep (ES, 27; castrated males) and Valaisian Black Nose Sheep (VS; 28; castrated males and females, 4:3), grazed for 9 weeks in paddocks allowing *ad libitum* forage intake on the following four pasture types: lowland control red/white clover-ryegrass ley, 0 % slope at 400 m a.s.l. (C), alpine *Crepido aurea-Festucetum* pasture, 0-5 % slope at 1950 m a.s.l. (F), alpine *Geo montani-Nardetum* pasture, >40 % slope at 2250 m a.s.l. (G) and alpine *Seslerio-Caricetum sempervirentis* pasture, >40 % slope at 2150 m a.s.l. (S). The initial live weights were 32.7±3.5 kg and 30.8±3.0 kg and ages were 26±2 and 18±7 weeks for ES and VS, respectively. Animals were balanced for gender, weight and age. Before and after the experimental period, markers of bone formation (bAP, bone-specific alkaline phosphatase) and bone resorption (ICTP, cross-linked carboxyterminal telopeptide of type I collagen) were analysed in blood samples and bone mineral density (BMD), bone mineral content (BMC) and cortical thickness (CT) measurements were conducted with peripheral quantitative computer tomography (pQCT) in the middle of the diaphysis on the left metatarsus. After slaughtering, maximum force (MF) to break the left metatarsus in the middle of the diaphysis was analyzed. Concentrations of Calcium (Ca) were analysed with COBAS MIRA[®] Autoanalyser in one pooled forage samples per pasture type. Data was subjected to analysis of variance considering pasture type, breed and the interaction of both. Furthermore, Pearson correlation coefficients were computed.

Results Concentrations (% DM) of Ca in forage samples were 0.54, 0.97, 0.50 and 1.62 for the pasture types C, F, G and S respectively, indicating a large variation in Ca supply of the lambs on the different pasture types. A trend ($P=0.054$) was found for a breed × pasture type interaction of BMD development. BMD increased during the experiment for ES by 101.6, 15.5, 24.8 and 48.1 mg/cm³ and for VS by 40.9, 97.5, 168.6 and 50.9 mg/cm³ on the pasture types C, F, G and S, respectively. The BMC increased during the experimental period for ES by 32.3 and 6.5 mg/cm and for VS by 5.0 and 33.8 mg/cm on C and F and even decreased for ES by 2.9 and 10.8 mg/cm and for VS by 22.8 and 10.6 on G and S, respectively. This results in pasture type differences ($P<0.001$), as well as a breed × pasture type interaction ($P<0.001$) for BMC development. The CT increased across all pasture types for both breeds during the experiment, but again (like for BMD) the increase in CT was higher for ES on C and for VS on the three alpine pastures (F, G and S), resulting in a significant breed × pasture type interaction ($P=0.02$). Further, a significant breed × pasture type interaction ($P=0.004$) as well as a significant breed difference ($P=0.001$) were found for MF to break the metatarsi in its middle. Metatarsi of VS were less stable than those of ES on the pasture types C, G and S while this was opposite for F. The pQCT measurements at the end of the experiment showed interestingly similar pattern for BMC, MF, BMD and CT although the development of those parameters during the experiment followed different ways. Close correlations were found between MF and BMC ($R^2=0.93$; $P<0.001$) and between CT and BMD ($r=0.94$; $P<0.001$). Between BMC and BMD a significant correlation only developed during the experiment, with $r=0.80$ ($P<0.001$) at the end compared to $r=0.25$ ($P=0.064$) at the beginning of the experiment. Calculation of modelling/remodelling rate (RMR) by dividing bAP through ICTP on both sampling dates gave a punctual impression of the bone metabolism of the lambs at those two specific dates. Significant pasture type differences ($P<0.001$) were found for RMR, as values decreased slightly during the experiment for both breeds on C and more clearly on G, whereas the decrease was remarkably stronger on both pasture types for ES than VS. This indicates that bone resorption outweighed bone formation processes, which corresponds with low Ca concentrations on C and G. Additionally, ES appears to be more sensitive in bone marker responds than VS. The RMR increased for both breeds during the experiment clearly on F and moderately on S, pointing towards facilitated bone formation processes on those pastures that had higher Ca concentration in the forage (F and S) compared to low Ca input pastures.

Conclusion Breed × pasture type interactions were found for BMC and CT development and also a trend for interactions for BMD development, indicating a different adaption of the two breeds to mountain pastures. The VS profit more from high alpine grazing compared to lowland grazing in terms of CT and BMD development while this was opposite for ES. The differences in alpine pastures characteristics, especially in slope, nutritional supply and also the individual movement behaviour of the lambs appear to influence BMC and RMR development throughout alpine grazing. Pasture types, including all characteristics mentioned, do have an important effect on bone metabolism of lambs and are tolerated by breeds to a different extent.

References

- Bass, L.S., Eser, P. and Daly, R. 2005. *Journal of Musculoskeletal and Neuronal Interactions*. 5(3): 239-254
Liesegang, A., Kaulfers, C., Lauff K. *et al.* 2010. *Bone*. 46, 74